

◆STEP 2: Emergency Contacts◆

1. Activate EMS and share relevant assessment and treatment information

2. Dr. _____ Phone Number: _____

3. Parent _____ Phone Number(s) _____

3. Emergency contacts: Name/Relationship _____ Phone Number(s) _____

a. _____ 1) _____ 2) _____

b. _____ 1) _____ 2) _____

Trained Staff:

1. _____ Room _____

2. _____ Room _____


3. _____ Room _____

EpiPen® and EpiPen® Jr. Directions

- Pull off gray activation cap




- Hold black tip near outer thigh (always apply to thigh)



- Swing and jab firmly into outer thigh until Auto-Injector mechanism functions.
- Hold in place and count to 10
- Remove the EpiPen® unit and massage the injection area for 10 seconds

Auvi-Q Directions

- Pull off outer cover/red activation cap
- Follow voice prompts for injection



Twinject® 0.3 mg and Twinject® 0.15 mg Directions

- Remove caps labeled “1” and “2”
- Place rounded tip against outer thigh, press down hard until needle penetrates
- Hold for 10 seconds, then remove



Second Dose Administration:
If symptoms don't improve after 10 minutes, administer second dose:

- Unscrew rounded tip. Pull syringe from barrel by holding blue collar at needle base
- Slide yellow collar off plunger
- Put needle into thigh through skin, push plunger down all the way, and remove

